arXiv:hep-th/9211137v1 30 Nov 1992

DILOGARITHM IDENTITIES AND SPECTRA IN CONFORMAL FIELD THEORY

ANATOL N. KIRILLOV
Isaac Newton Institute for Mathematical Sciences,
20 Clarkson Road, Cambridge, CB3 OEH, U.K.
and
Steklov Mathematical Institute,
Fontanka 27, St.Petersburg, 191011, Russia

ABSTRACT

We prove new identities between the values of Rogers dilogarithm function and describe a connection between these identities and spectra in conformal field theory.

The dilogarithm function defined for $0 \leq z \leq 1$ by

$$Li_2(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^2} = -\int_0^z \frac{\log(1-t)}{t} dt,$$

is one of the lesser transcendental functions. Nonetheless, it has many intriguing properties and has appeared in various branches of mathematics and physics. Here we mention only the appearences of dilogarithm function in number theory (the study of asymptotic behaviour of partitious [RS]; the values of ζ-functions in some special points [Za]) in algebraic K-theory (A.Beilinson, S.Bloch, A.Groncharov,...).

In physics, the dilogarithm appears at first from a calculation of magnetic susceptibility in the XXZ model at small magnetic field [KR]. More recently [Z], the dilogarithm identities (through the Thermodynamic Bethe Ansatz (TBA)) through appear in the context of investigation of UV limit or the critical behaviour of integrable 2-dimensional quantum field theories and lattice models [Z], [DR], [KP],...). One aim of this paper is to prove some new identities between the values of Rogers dilogarithm function and to show that any rational number may be obtained as the value of some dilogarithm sum. More detailes and further results will appear in [Ki2].

§1. Definition and the basic properties of Rogers dilogarithm.

Let us remind the definition of the Rogers dilogarithm function $L(x)$ for $x \in (0,1)$

$$L(x) = -\frac{1}{2}\int_0^x \left[\frac{\log(1-x)}{x} + \frac{\log x}{1-x}\right] dx = \sum_{n=1}^{\infty} \frac{x^n}{n^2} + \frac{1}{2}\log x \cdot \log(1-x). \tag{1.1}$$

The following two classical results (see e.g. [Le], [GM], [Ki]) contain the basic properties of the function $L(x)$.

Theorem A. The function $L(x) \in C^{\infty}((0,1))$ and satisfies the following functional equations

$$1.\ \ L(x) + L(1-x) = \frac{\pi^2}{6}, \quad 0 < x < 1, \tag{1.2}$$

$$2.\ \ L(x) + L(y) = L(xy) + L\left(\frac{x(1-y)}{1-xy}\right) + L\left(\frac{y(1-x)}{1-xy}\right), \tag{1.3}$$

where $0 < x,\ y < 1$.

Theorem B. Let$f(x)$ be a function of class $C^3((0,1))$ and satisfies the relations (1.2) and (1.3). Then we have

$$f(x) = \mathrm{const} \cdot L(x)$$

We continue the function $L(x)$ on all real axis $\mathbf{R} = \mathbf{R}^1 \cup \{\pm\infty\}$ by the following rules

$$L(x) = \frac{\pi^2}{3} - L(x^{-1}), \quad \text{if} \quad x > 1, \tag{1.4}$$

$$L(x) = L\left(\frac{1}{1-x}\right) - \frac{\pi^2}{6}, \quad \text{if} \quad x < 0, \tag{1.5}$$

$$L(0) = 0, \quad L(1) = \frac{\pi^2}{6}, \quad L(+\infty) = \frac{\pi^2}{3}, \quad L(-\infty) = -\frac{\pi^2}{6}.$$

The present work will concern with relations between the values of the Rogers dilogarithm function at certain algebraic numbers. More exactly, let us consider an abelian subgroup W in the field of rational numbers $\mathbf{Q}$

$$W = \left\{\sum_i \frac{n_i L(\alpha_i)}{L(1)} \mid n_i \in \mathbf{Z}, \quad \alpha_i \in \overline{\mathbf{Q}} \cap \mathbf{R} \ \text{ for all } \ i\right\} \cap \mathbf{Q}. \tag{1.6}$$

According to a conjecture of W. Nahm [Na] the abelian group W "coincides" with the spectra in rational conformal field theory. Thus it seems very interesting task to obtain more explicit description of the group W (e.g. to find a system of generators for W) and also to connect already known results about the spectra in conformal field theory (see e.g. [BPZ], [GKO], [FF], [FQS], [Ka], [KW], [KP]) with suitable elements in W. One of our main results of the present paper allows to describe some part of a system of generators for abelian group W. As a corollary, we will show that the spectra of unitary minimal models [BPZ] and some others are really contained in W.

§2. Basic identities and conformal weights.

In this section we present our main results dealing with a computation of the following dilogarithm sum

$$\sum_{k=1}^{n-1}\sum_{m=1}^{r} L\left(\frac{\sin k\varphi\cdot\sin(n-k)\varphi}{\sin(m+k)\varphi\cdot\sin(m+n-k)\varphi}\right) := \frac{\pi^2}{6}s(j,n,r), \tag{2.1}$$

where $\varphi=\dfrac{(j+1)\pi}{n+r}$, $0\le j\le n+r-2$.

It is clear that $s(j,n,r)=s(n+r-2-j,n,r)$, so we will assume in sequel that $0\le 2j\le n+r-2$.

The dilogarithm sum (2.1) corresponds to the Lie algebra of type A_{n-1}. The case $j=0$ was considered in our previous paper [Ki], where it was proved that $s(0,n,r)=\dfrac{(n^2-1)r}{n+r}$. It was stated in [Ki] that this number coincides with the central charge of the $SU(n)$ level r WZNW model. Before formulating our result about computation of the sum $s(j,n,r)$ let us remind the definition of Bernoulli polynomials. They are defined by the generating function

$$\frac{te^{xt}}{e^t-1}=\sum_{n=0}^{\infty}B_n(x)\frac{t^n}{n!},\quad |t|<2\pi.$$

We use also modified Bernoulli polynomials

$$\overline{B}_n(x)=B_n(\{x\}),\quad \text{where}\ \ \{x\}=x-[x]$$

be a fractional part of $x\in\mathbf{R}$. It is well-known that

$$\overline{B}_{2n}(x)=(-1)^n\frac{2(2n)!}{(2\pi)^{2n}}\sum_{k=1}^{\infty}\frac{\cos 2k\pi x}{k^{2n}}, \tag{2.2}$$

$$\overline{B}_{2n+1}(x)=(-1)^{n-1}\frac{2(2n+1)!}{(2\pi)^{2n+1}}\sum_{k=1}^{\infty}\frac{\sin 2k\pi x}{k^{2n+1}}.$$

Theorem 2.1. We have

$$s(j,n,r)=6(r+n)\sum_{k=0}^{[\frac{n-1}{2}]}\left\{\frac{1}{6}-\overline{B}_2\Big((n-1-2k)\theta\Big)\right\}-\frac{1}{4}\Big\{2n^2+1+3(-1)^n\Big\}, \tag{2.3}$$

where $\theta=\dfrac{j+1}{r+n}$ and g.c.d.$(j+1,r+n)=1$.

Theorem 2.2. (level-rank duality)

$$s(j,n,r)+s(j,r,n)=nr-1. \tag{2.4}$$

Corollary 2.3. We have

$$s(j,n,r)=c_r^{(n)}-24h_j^{(r,n)}+6\cdot\mathbf{Z}_+, \tag{2.5}$$

where

$$c_r^{(n)}=\frac{(n^2-1)r}{n+r},\quad h_j^{(r,n)}=\frac{n(n^2-1)}{24}\cdot\frac{j(j+2)}{r+n},\quad 0\le j\le r+n-2, \tag{2.6}$$

are the central charge and conformal dimensions of the $SU(n)$ level r WZNW primary fields, respectively.

Proof. Let us remind that

$$B_2(x)=x^2-x+\frac{1}{6}\quad\text{and}\quad\overline{B}_2(x)=B_2(\{x\}).$$

Thus,

$$\begin{aligned}
&6(r+n)\sum_{k=0}^{[\frac{n-1}{2}]}\left(\frac{1}{6}-\overline{B}_2\Big((n-1-2k)\theta\Big)\right)=\\
&=6(r+n)\sum_{k=0}^{[\frac{n-1}{2}]}(n-1-2k)\theta-6(r+n)\sum_{k=0}^{[\frac{n-1}{2}]}(n-1-2k)^2\theta^2+\\
&+6(r+n)\sum_{k=0}^{[\frac{n-1}{2}]}\Big[(n-1-2k)\theta\Big]\left((n-1-2k)\theta-1+\Big\{(n-1-2k)\theta\Big\}\right):=\\
&=6\Sigma_1-6\Sigma_2+6\Sigma_3.
\end{aligned}$$

Now if we take $\theta=\dfrac{j+1}{r+n}$, then it is clear that $\Sigma_3\in\mathbf{Z}_+$. In order to compute Σ_1 and Σ_2 we use the following summation formulae

$$\begin{aligned}
&\sum_{k=0}^{[\frac{n-1}{2}]}(n-1-2k)=\frac{2n^2-1+(-1)^n}{8}=\left[\frac{n}{2}\right]\cdot\left[\frac{n+1}{2}\right],\\
&\sum_{k=0}^{[\frac{n-1}{2}]}(n-1-2k)^2=\frac{n(n^2-1)}{6}.
\end{aligned}$$

Consequently $\quad s(j,n,r)=$

$$\begin{aligned}
&=\frac{3(2n^2-1+(-1)^n)(j+1)}{4}-\frac{n(n^2-1)(j+1)^2}{r+n}-\frac{2n^2+1+3(-1)^n}{4}+6\Sigma_3=\\
&=\frac{(n^2-1)r}{r+n}-\frac{n(n^2-1)j(j+2)}{r+n}+6j\left[\frac{n}{2}\right]\cdot\left[\frac{n+1}{2}\right]+6\Sigma_3.
\end{aligned}$$

■

For small values of j we may compute the sum in (2.3) and thus to find corresponding positive integer in (2.5).

Corollary 2.4.

i) if $j \leq r$, then

$$s(j,2,r) = c_r^{(2)} - 24h_j^{(r,2)} + 6j = \frac{3r}{r+2} + 6\,\frac{j(r-j)}{r+2}. \tag{2.7}$$

ii) if $2j \leq r+1$, then

$$s(j,3,r) = \frac{8r}{r+3} - 24\,\frac{j(j+2)}{r+3} + 12j. \tag{2.8}$$

iii) if $(n-1)j < r+1$, then

$$s(j,n,r) = c_r^{(n)} - 24h_j^{(r,n)} + 6j \cdot \left[\frac{n}{2}\right] \cdot \left[\frac{n+1}{2}\right]. \tag{2.9}$$

Proof. An assumption $(n-1)j < r+1$ is equivalent to a condition $\dfrac{(n-1)(j+1)}{n+r} < 1$. So the term Σ_3 (see a proof of Corollary 2.3) is equal to zero. ■

It seems interesting to find a meaning of the positive integer in (2.5). Now we want to find a "dilogarithm interpretation" of the central charges and conformal dimensions for some well-known conformal models.

Corollary 2.5. We have

$$s(j_1,2,k)+s(0,2,1)-s(j_2,2,k+1) = c_k - 24h_{j_1+1,j_2+1} + 6(j_1-j_2)(j_1-j_2+1), \tag{2.10}$$

where

$$c_k = 1 - \frac{6}{(k+2)(k+3)}, \quad h_{r,s}^{(k)} = \frac{[(k+3)r-(k+2)s]^2-1}{4(k+2)(k+3)} \tag{2.11}$$

are the central charge and conformal dimensions of the primary fields for unitary minimal conformal models [BPZ], [Ka], [FQS].

Corollary 2.6.

$$s(j_1,n,k) + s(0,n,1) - s(j_2,n,k+1) = c_{k,n} - 24h_{j_1+1,j_2+1}^{(n)} + 12\mathbf{Z}_+, \tag{2.12}$$

where

$$\begin{aligned} c_{k,n} &= (n-1)\left\{1 - \frac{n(n+1)}{(k+n)(k+n+1)}\right\}, \\ h_{r,s}^{(n)}(k) &= \frac{n(n^2-1)}{24} \cdot \frac{[(k+n+1)r-(k+n)s]^2-1}{(k+n)(k+n+1)} \end{aligned} \tag{2.13}$$

are the central charge and conformal dimensions of the primary fields for W_n models [Bi].

Finally we give a "dilogarithm interpretation" for the central charge and conformal weights of restricted solid-on-solid (RSOS) lattice models and their fusion hierarchies [KP].

Corollary 2.7. We have

$$\begin{aligned} & s(l,2,N)+s(N-1,2,N-2)-s(m-1,2,N-2)= \\ & =c+1-24\Delta+6(l-|m|), \quad m\in\mathbf{Z}, \quad 0\le l\le N, \end{aligned} \tag{2.14}$$

where

$$c=\frac{2(N-1)}{N+2} \quad \text{and} \quad \Delta=\frac{l(l+2)}{4(N+2)}-\frac{m^2}{4N} \tag{2.15}$$

are the central charge and conformal weights of $\mathbf{Z}_N$ parafermion theories [FZ]. The members of (2.15) may be also realized as the central charge and conformal weights of fusion $N+1$-state RSOS(p,p) lattice models [DJKMO], [BR] on the regime I/II critical line. Note that physical constraints

$$|m|\le l, \quad m\equiv l \ (\mathrm{mod}\ 2)$$

for value of m in (2.15) are equivalent to a condition that "remainder term" in (2.14), namely, $6(l-|m|)$, must be belongs to $12\mathbf{Z}_+$.

Corollary 2.8. Let us fix the positive integers k, $p=1,2,\ldots$ (the fusion level), j_1 and j_2 such that $0\le j_1\le k$, $0\le j_2\le k+l$.

Let us put $r_0=p\left\{\dfrac{j_1-j_2}{p}\right\}$. Then we have

$$\begin{aligned} & s(j_1,2,k)+s(r_0,2,p)-s(j_2,2,k+p)= \\ & =c-24\Delta+12\ \frac{(j_1-j_2)(p+j_1-j_2)+r_0(p-r_0)}{2p}, \end{aligned} \tag{2.16}$$

where

$$c=\frac{3p}{p+2}\left(1-\frac{2(p+2)}{(k+2)(k+p+2)}\right), \tag{2.17}$$

$$\Delta=\frac{[(k+p+2)(j_1+1)-(k+2)(j_2+1)]^2-p^2}{4p(k+2)(k+p+2)}+\frac{r_0(p-r_0)}{2p(p+2)}$$

are the central charge and conformal weights of the fusion $(k+p+1)$-state RSOS(p,p) latice models [KP] on the regime III/IV critical line. It is easy to see that "remainder term" in (2.16) belongs to $12\mathbf{Z}_+$. Note also that the fusion RSOS(p,q) lattice models, obtained by fusing $p\times q$ blocks of face weights together, are related to coset conformal fields theories obtained by the Goddard-Kent-Olive (GKO) construction [GKO]. Namely, c and Δ in (2.17) are the central charge and conformal dimensions of conformal field theory, which corresponds to the coset pair [GKO]

$$\begin{array}{cccccc} & A_1 & \oplus & A_1 & \supset & A_1 \\ \text{levels} & k & & p & & k+p \end{array}$$

§3. A_1-type dilogarithm identities.

As is well-known [Le], the Rogers dilogarithm function $L(x)$ admits a continuation on all complex plane $\mathbf{C}$. Follow [Le], [KR] we define a function

$$L(x,\theta):=-\frac{1}{2}\int_0^x \frac{\log(1-2x\cos\theta+x^2)}{x}dx+\frac{1}{4}\log|x|\cdot\log(1-2x\cos\theta+x^2)= \\ =ReL(xe^{i\theta}),\quad x,\theta\in\mathbf{R} \tag{3.1}$$

Our proof of Theorem 2.1 is based on a study of properties of the function $L(x,\theta)$.

Proposition 3.1. For all real φ, θ we have

$$L\Big(\Big(\frac{\sin\theta}{\sin\varphi}\Big)^2\Big)=\pi^2\Big\{\overline{B}_2\left(\frac{\theta+\varphi}{\pi}\right)-\overline{B}_2\left(\frac{\varphi}{\pi}\right)-\overline{B}_2\left(\frac{\theta}{\pi}\right)+\frac{1}{6}\Big\}+ \\ +2L\Big(-\frac{\sin(\varphi-\theta)}{\sin\theta},\varphi\Big)-2L\Big(-\frac{\sin\varphi}{\sin\theta},\varphi+\theta\Big). \tag{3.2}$$

Before proving a Proposition 3.1 let us give the others useful properties of function (3.1) (compare with [Le]).

Lemma 3.2.

(i) $L(x,0)=L(x),\quad L(-x,\varphi)=L(-x,\pi-\varphi)$ (3.3)

(ii) $L(x,\varphi)=L(x,2\pi k\pm\varphi),\quad k\in\mathbf{Z}$ (3.4)

(iii) $L(-1,\varphi)=\pi^2\overline{B}_2\left(\frac{\varphi}{2\pi}+\frac{1}{2}\right),$

$L(1,\varphi)=\pi^2\overline{B}_2\left(\frac{\varphi}{2\pi}\right)$ (3.5)

(iv) $L(x,\varphi)+L(x^{-1},\varphi)=2\pi^2\overline{B}_2\left(\frac{\varphi}{2\pi}\right),\quad x>0$

$L(-x,\varphi)+L(-x^{-1},\varphi)=2\pi^2\overline{B}_2\left(\frac{\varphi}{2\pi}+\frac{1}{2}\right),\quad x<0$ (3.6)

(v) $L(0,\varphi)=0,\quad L(+\infty,\varphi)=2\pi^2\overline{B}_2\left(\frac{\varphi}{2\pi}\right),$

$L(-\infty,\varphi)=2\pi^2\overline{B}_2\left(\frac{\varphi+\pi}{2\pi}\right)$ (3.7)

(vi) $L(2\cos\varphi,\varphi)=\pi^2\Big\{\overline{B}_2\left(\frac{\varphi}{\pi}\right)+\frac{1}{12}\Big\}$ (3.8)

(vii) $L(x^n,n\varphi)=n\sum_{k=0}^{n-1}L\Big(x,\ \varphi+\frac{2k\pi}{n}\Big),\quad x\in\mathbf{R}_+,$

$L(x^n)=n\sum_{k=0}^{n-1}L\left(x\cdot\exp\frac{2k\pi i}{n}\right),\quad x\in(0,1).$ (3.9)

More generally (Rogers' identity [Ro])

$$L(1-y^n)=\sum_{k=1}^n\sum_{l=1}^n\Big[L(\lambda_k/\lambda_l)-L(x_k\lambda_l)\Big],$$

where $\{x_k\}_{k=1}^n$ are the roots of the equation

$$1-y^n=\prod_{k=1}^n(1-\lambda_k x).$$

Proof of the Theorem 2.1 for the case $n=2$. If we substitute $\theta=m\varphi$ in (3.2) then obtain

$$L\left(\left(\frac{\sin m\varphi}{\sin\varphi}\right)^2\right)=\pi^2\left\{\overline{B}_2\left(\frac{(m+1)\varphi}{\pi}\right)-\overline{B}_2\left(\frac{m\varphi}{\pi}\right)-\overline{B}_2\left(\frac{\varphi}{\pi}\right)+\frac{1}{6}\right\}+$$
$$+2L\left(-\frac{\sin((m-1)\varphi)}{\sin\varphi},m\varphi\right)-2L\left(-\frac{\sin m\varphi}{\sin\varphi},(m+1)\varphi\right) \tag{3.10}$$

Futher let us introduce notations

$$f_m(\varphi):=1-\frac{Q_{m-1}(\varphi)Q_{m+1}(\varphi)}{Q_m^2(\varphi)}=\frac{1}{Q_m^2(\varphi)}.$$

Then using (3.10) we find

$$\sum_{m=1}^r L(f_m(\varphi))=-2\left\{L\Big(-Q_r(\varphi),(r+2)\varphi\Big)-\frac{\pi^2}{6}\right\}+ \tag{3.11}$$
$$+\pi^2\left\{\overline{B}_2\left(\frac{(r+2)\varphi}{\pi}\right)-\frac{1}{6}\right\}+(r+2)\pi^2\left\{\frac{1}{6}-\overline{B}_2\left(\frac{\varphi}{\pi}\right)\right\}-\frac{\pi^2}{2}.$$

Now let us put $\varphi=\dfrac{(j+1)\pi}{r+2}$, $0\le j\le r+1$. Then $Q_r(\varphi)=(-1)^j$ and it is clear from (3.3) and (3.4) that

$$L\Big((-1)^{j+1},\ (j+1)\pi\Big)=L(1)=\frac{\pi^2}{6}.$$

■

Note that polynomials $Q_m:=Q_m(\varphi)$ satisfy the following recurrence relation

$$Q_m^2=Q_{m-1}Q_{m+1}+1,\quad Q_0\equiv 1,\quad m\ge 1,$$

where as the polynomials $y_m:=y_m(\varphi)=Q_{m-1}(\varphi)\cdot Q_{m+1}(\varphi)$ satisfy the following one

$$y_m^2=(1+y_{m-1})(1+y_{m+1}),\quad y_0\equiv 0,\quad m\ge 1.$$

Now we propose a generalisation of (3.12). Given a rational number p and decomposition of p into the continued fraction

$$p=[b_r,b_{r-1},\dots,b_1,b_0]=b_r+\cfrac{1}{b_{r-1}+\cfrac{1}{\cdots+\cfrac{1}{b_1+\cfrac{1}{b_0}}}}. \tag{3.12}$$

We will assume that $b_i > 0$ if $0 \leq i < r$ and $b_r \in \mathbf{Z}$. Using the decomposition (3.12) we define the set of integers y_i and m_i:

$$\begin{aligned} &y_{-1} = 0, \quad y_0 = 1, \quad y_1 = b_0, \dots, y_{i+1} = y_{i-1} + b_i y_i, \quad 0 \leq i \leq r, \\ &m_0 = 0, \quad m_1 = b_0, \quad m_{i+1} = |b_i| + m_i, \quad 0 \leq i \leq r. \end{aligned} \tag{3.13}$$

The following sequences of integers were first introduced by Takahashi and Suzuki [TS]: $r(j) = i$, if $m_i \leq j < m_{i+1}$, $0 \leq i \leq r$,

$$n_j = y_{i-1} + (j - m_i) y_i, \quad \text{if} \quad m_i \leq j < m_{i+1} + \delta_{i,r}, \quad 0 \leq i \leq r.$$

Finally we define a dilogarithm sum of "fractional level p":

$$\sum_{j=1}^{m_{r+1}} (-1)^{r(j)} L\left(\left(\frac{\sin y_{r(j)}\theta}{\sin(n_j + y_{r(j)})\theta} \right)^2 \right) := (-1)^r \frac{\pi^2}{6} s(k,2,p), \tag{3.14}$$

where $\theta = \dfrac{(k+1)\pi}{y_{r+1} + 2y_r}$

Proposition 3.3. We have

$$s(k,2,p) = \frac{3p}{p+2} - 6\frac{k(k+2)}{p+2} + 6\mathbf{Z}. \tag{3.15}$$

Conjecture 3.4. For all positive $p \in \mathbf{Q}$ the remainder term in (3.15) lies in $6\mathbf{Z}_+$.

Corollary 3.5. Let us fix the positive integers $l = 1, 2, 3, \dots$ (the fusion level), $p > q$, j_1 and j_2. Then

$$s\left(j_1, 2, \frac{ql}{p-q} - 2\right) + s(r_0, 2, l) - s\left(j_2, 2, \frac{pl}{p-q} - 2\right) = c - 24\Delta + 6\mathbf{Z}, \tag{3.16}$$

where $r_0 = l \cdot \left\{ \dfrac{j_1 - j_2}{l} \right\}$ and

$$\begin{aligned} c &= \frac{3l}{l+2}\left(1 - \frac{2(l+2)(p-q)^2}{l^2 pq}\right), \\ \Delta &= \frac{[p(j_1+1) - q(j_2+1)]^2 - (p-q)^2}{4lpq} + \frac{r_0(l - r_0)}{2l(l+2)} \end{aligned} \tag{3.17}$$

are the central charge and conformal dimensions of RCFT, which corresponds to the coset pair [GKO]

$$\begin{array}{ccccc} & A_1 & \oplus\ A_1 & \supset & A_1 \\ \text{levels} & \dfrac{ql}{p-q} - 2 & l & & \dfrac{pl}{p-q} - 2 \end{array}$$

Acknowledgements. Most of this work was done during a visit to the Isaac Newton Institute for Mathematical Sciences. I thank W.Nahm, E.K.Sklyanin, F.Ravanini and E.Corrigan for very helpfull discussions.

Riferences.

[Bi] A.Bilal, Nucl.Phys., B330, (1990), 399.

[BPZ] A.Belavin, A.Polyakov, A.Zamolodchikov, J.Stat.Phys., 34, (1984), 763; Nucl. Phys., B241, (1984), 333.

[BR] V.V.Bazhanov, N.Yu.Reshetikhin, Prog.Theor.Phys., 102, (1990), suppl.

[CR] P.Christe, F.Ravanini, $G_N \otimes G_L/G_{N+L}$ conformal field theories and their modular invariant partition functions, Int.J.Mod.Phys., A4, (1989), 897.

[DR] P.Dorey, F.Ravanini, Staircase model from affine Toda field theory, prepr. SPhT/92-065.

[FL] V.A.Fateev, S.Lukyanov, Int.J.Mod.Phys.,A3, (1988), 507.

[FQS] D.Friedan, Z.Qiu, S.Shenker, Phys.Rev.Lett., 52, (1984), 1575.

[GKO] P.Goddard, A.Kent, D.Olive, Phys.Lett., B152, (1985), 105.

[GM] I.M.Gelfand, R.D.MacPherson, Advan. in Math., 44, (1982), 279.

[Ka] V.G.Kac, Infinite dimensional Lie algebras, Cambridge Univ.Press. 1990.

[Ki1] A.N.Kirillov, Zap.Nauch.Semin. LOMI, 164, (1987), 121.

[Ki2] A.N.Kirillov, Dilogarithm identities, partitions and spectra in conformal field theory, in preparation.

[KN1] A.Kuniba, T.Nakanishi, Level-rank duality in fusion RSOS models, prepr. 1989.

[KN2] A.Kuniba, T.Nakanishi, Spectra in conformal field theories from Rodgers dilogarithm, prepr.SMS-042-92, 1992.

[KP] A.Klumper, P.Pearce, Physica, A183, (1992), 304.

[KR] A.N.Kirillov, N.Yu.Reshetikhin, Zap.Nauch.Semin. LOMI, 160, (1987), 211.

[Ku] A.Kuniba, Thermodynamics of the $U_q(X_r^{(1)})$ Bethe ansatz system with q a root of Unity, ANU, prepr. SMS-098-91.

[KW] V.Kac, M.Wakimoto, Adv.Math., 70, (1988), 156.

[Le] L.Lewin, Polylogarithms and associated functions, (North-Holland, 1981).

[MSW] P.Mathieu. D.Senechal, M.Walton, Field identification in nonunitary diagonal cosets, prepr. LETH-PHY-9/91.

[Na] W.Nahm, Dilogarithm and W-algebras, Talk given at the Isaac Newton Inst., Cambridge, Sept. 1992.

[NRT] W.Nahm, A.Recknagel, M.Terhoeven, Dilogarithm identities in conformal field theory, BONN-prepr., hepth 9211034.

[Ro] L.J.Rogers, Proc.London Math Soc., 4, (1907), 169.

[RS] B.Richmond, G.Szekeres, J.Austral.Math.Soc., A31, (1981), 362.

[SA] H.Saleur, D.Altshuler, Nucl.Phys., B354, (1991), 579.

[TS] M.Takahashi, M.Suzuki, Prog.Theor.Phys., 48, (1972), 2187.

[Z] Al.B.Zamolodchikov, Nucl.Phys., B342, (1990), 695.

[Za] D.Zagier, The remarkable dilogarithm (Number theory and related topics. Papers presented at the Ramanujan colloquium, Bombay, 1988, TIFR).